

Functional Domains of the Epithelial Sodium Channel

Johannes Loffing* and Laurent Schild[†]

*University of Fribourg, Department of Medicine, Unit of Anatomy, Fribourg, Switzerland; and [†]Department de Pharmacologie & Toxicologie, Faculté de Biologie et Médecine, Université de Lausanne, Lausanne, Switzerland

The epithelial sodium channel (ENaC) plays an important role in ion transport of many organ systems. In kidney, colon, and sweat gland, ENaC-mediated trans-epithelial sodium transport helps to adjust sodium excretion in the urine, feces, and sweat to fulfill homeostatic requirements. In lung and airways, ENaC activity is important for alveolar liquid clearance and regulation of mucous fluidity. In the taste buds of the tongue, ENaC is likely involved in salt tasting, whereas ENaC expression in the eye and inner ear may help to control the ionic composition of the aqueous humor and the endolymph, respectively. These diverse functions of ENaC (1) require intricate molecular regulatory systems that precisely adapt ENaC function to constantly changing electrolyte transport requirements. The best studied regulatory system is the aldosterone-dependent regulation of ENaC in the kidney, which is of major importance for the maintenance of whole-body sodium homeostasis and for BP control (2). This review focuses exclusively on the role of ENaC in renal sodium handling. Distribution and aldosterone-dependent regulation of ENaC in the kidney *in vivo* is discussed briefly, and the functional domains of ENaC that are important for the regulation of the channel under physiologic and pathophysiologic conditions are highlighted.

ENaC and the Aldosterone-Sensitive Distal Nephron

ENaC is the major apical sodium entry pathway present in the aldosterone-sensitive distal nephron (ASDN). The ENaC-mediated sodium transport across the apical membrane of tubule cells is electrogenic, depolarizing this membrane and favoring K⁺ secretion *via* apical K⁺ channels such as ROMK. Moreover, sodium transport across ENaC may osmotically drive transepithelial water transport *via* vasopressin-dependent apical water channels (*i.e.*, aquaporin-2). As such ENaC in the

ASDN plays a pivotal role in the final adjustment of renal sodium, potassium, and water excretion. The importance of ENaC for sodium, potassium, and fluid homeostasis is emphasized by the observation that ENaC gain-of-function mutations or loss-of-function mutations lead, respectively, to extracellular volume expansion (Liddle's syndrome or pseudohyperaldosteronism) or to renal salt-wasting syndromes (pseudohypoaldosteronism type 1) associated with alterations in potassium homeostasis (2,3).

Morphologic and functional studies on rodent and human kidneys (4–6) indicated that at least three successive tubule portions—the late portion of the distal convoluted tubule (DCT), the connecting tubule (CNT), and the collecting duct (CD)—contribute to the ASDN. Although these segments have distinct structural and functional features (6), they share in common the expression of ENaC, the mineralocorticoid receptor (MR), and the 11- β hydroxysteroid dehydrogenase type 2 proteins (5). The latter confers mineralocorticoid-selectivity to the MR by rapid metabolism of circulating glucocorticoids. It is uncertain whether the early DCT also belongs to the ASDN. The early DCT weakly expresses the MR but lacks detectable 11- β hydroxysteroid dehydrogenase type 2 (7). In the kidney, aldosterone increases the protein expression of the DCT-specific NaCl co-transporter that is sensitive to thiazides (NaCl co-transporter or thiazide-sensitive co-transporter) (8,9). Consistently, aldosterone has been shown to stimulate electroneutral Na⁺ transport in the DCT (10). It needs to be determined whether these stimulatory effects occur along the entire DCT or only in the late DCT.

The subcellular localization of ENaC along the axis of the ASDN changes drastically with the elevation of plasma aldosterone levels in response to changes in the sodium diet (11,12). In rodents that are kept under a high dietary sodium intake with low plasma aldosterone levels, ENaC subunits are barely detectable at the luminal membrane and are found almost exclusively at intracellular sites, the identity of which remain to be identified. On a standard dietary sodium intake (European laboratory diet) with moderate plasma aldosterone levels, ENaC subunits are traceable at the luminal membrane of late DCT and early CNT. However, in segments farther downstream (late CNT and CD), particularly β - and γ -ENaC subunits remain almost exclusively localized at intracellular sites. Under a low dietary sodium intake with high plasma aldosterone

Address correspondence to: Dr. Johannes Loffing, University of Fribourg, Department of Medicine, Unit of Anatomy, Route Albert Gockel 1, 1700 Fribourg, Switzerland. Phone: +41-26-300 8527; Fax: +41-26-300-9733; E-mail: johannes.loffing@unifr.ch; Dr. Laurent Schild, Université de Lausanne, Department de Pharmacologie & Toxicologie, Faculté de Biologie et Médecine, rue du Bugnon 27, 1005 Lausanne, Switzerland. Phone: +41-21-692-5380; Fax: +41-21-692-5355; E-mail: laurent.schild@unil.ch

levels, ENaC becomes detectable in the luminal membrane along the late DCT, CNT, and CD (11,12). Nevertheless, the axial gradient for apical ENaC still prevails, and the apical localization of ENaC subunits is more prominent in early ASDN than in late ASDN (Figure 1). This immunohistochemically traceable axial gradient of apical ENaC localization is corroborated by a recent series of elegant patch-clamp studies (13). ASDN segments that were isolated from animals that were on a standard US laboratory diet exhibited no amiloride-sensitive currents at the single channel level, whereas CNT and CD that were isolated from rats with elevated plasma aldosterone levels revealed significant single-channel ENaC currents. In general, amiloride-sensitive currents decreased in the following order: CNT > initial CD > CD (13). These findings on apical ENaC localization and activity are consistent with previous studies on microperfused rat tubules (14,15) and on isolated rabbit tubules (16) that established several times higher sodium transport rates in early ASDN (DCT and CNT) than in ASDN segments farther downstream (CD), and also with studies that showed a progressive decrease of the basolateral Na-K-ATPase activity along the ASDN (17). Taken together, the data clearly suggest that the aldosterone-dependent adaptation of renal sodium excretion to dietary sodium intake occurs predomi-

nately in the early ASDN, whereas the late ASDN gets recruited only under high plasma aldosterone levels. The importance of the early ASDN *versus* late ASDN for the maintenance of sodium balance was recently highlighted by the development of a mouse model with targeted inactivation of α -ENaC exclusively in the CD. These mice survive well and are able to maintain sodium and potassium balance, even when challenged by salt restriction or potassium loading (18).

Functional Properties

Electrophysiologic investigations of ions transport first in the toad skin and in toad urinary bladder, then in isolated perfused distal nephrons have identified large amiloride-sensitive electrogenic Na^+ currents stimulated by aldosterone (19). The first recordings of amiloride-sensitive Na^+ currents at the single-channel level were obtained from principal cells of microdissected cortical CD and from cell lines and revealed the functional signature of the epithelial sodium channel (20,21). This channel was highly selective for Na^+ over K^+ ions ($P_{\text{Na}}/P_{\text{K}} > 100$); the single-channel conductance was 4 to 5 pS with Na^+ as the charge carrier and 9 to 10 pS with Li^+ ions. From patch-clamp recordings, ENaC usually shows long open and closed times. The open probability (P_o) of ENaC, however, is variable under similar physiologic conditions and range from ≤ 0.05 and > 0.95 ; this variation in P_o could reflect different gating modes of the channel (22). Switch between gating modes can be influenced by changes in the membrane voltage, although the response to voltage is relatively weak compared with the classical voltage-gated ion channels. Other variables have been proposed to affect ENaC gating and include intracellular pH and Ca^{2+} and hormones (23,24). High extracellular concentrations of Na^+ ions tend to inhibit ENaC activity, a phenomenon called self-inhibition (25). This fast inhibition of ENaC upon increasing extracellular Na^+ concentration is not related to changes in intracellular Na^+ concentration and is strongly dependent on the temperature.

From a pharmacologic point of view, ENaC is blocked by submicromolar concentrations of amiloride. Amiloride is a weak base, and the pH dependence of the block suggests that the ionized form of amiloride is efficient. Consistent with this notion, the amiloride block is voltage dependent, a characteristic that is commonly found for charged blockers binding in the channel pore within the transmembrane electric field (26). ENaC inhibition by amiloride is also dependent on the external concentration of Na^+ ions, and experiments in toad bladder are consistent with a competitive interaction between the permeating Na^+ ion and the blocker (27). These observations suggested overlapping binding sites for amiloride and Na^+ ions in the external pore vestibule of the channel.

The primary structure of ENaC was identified by expression cloning in *Xenopus* oocytes. ENaC is a heteromultimeric channel that is made of three homologous α , β , and γ subunits that share approximately 30% homology at the amino acid level (28,29). The expression of the $\alpha\beta\gamma$ -ENaC subunits recapitulates the functional and the pharmacologic properties of the channel (29). Predictions from the primary structure are consistent with the presence of two transmembrane α helices. The large loop




Figure 1. Apical localization of epithelial sodium channel (ENaC) along the aldosterone-sensitive distal nephron. On a high dietary sodium intake, ENaC subunits are almost absent from the cell surface and reside exclusively at intracellular sides in the aldosterone-sensitive distal nephron (ASDN) segments (the late distal convoluted tubule [DCT], the connecting tubule [CNT], and the collecting duct [CD]). Lowering dietary sodium intake progressively increases the cell surface abundance of ENaC starting in the early ASDN (*i.e.*, late DCT and early CNT) under moderate Na^+ intake and extending into the late ASDN under more severe Na^+ restriction. Shading of the tubules corresponds to the amount of apical ENaC and indicates the axial gradient of ENaC cell surface abundance along the ASDN. Illustration by Josh Gramling—Gramling Medical Illustration.

that separates the two transmembrane domains is facing the extracellular side as shown by the glycosylation pattern of the protein, although hydrophobic segments within this putative extracellular loop might be embedded in the membrane (see Figure 2) (30). According to these predictions, the N- and C-termini are facing the cytosolic side of the membrane. Most of the reports on the subunit stoichiometry of members of the ENaC/degenerin channel family agree with a four- subunit architecture, and in the case of ENaC, the channel is likely made of 2 α , 1 β , and 1 γ subunits (31–34). The issue of the subunit stoichiometry is not solved because ENaC channels with up to nine subunits have been proposed (35). From our knowledge of three-dimensional structure of different ion channels, such a nonameric structure, is unusual for highly selective ion channels. Such a high number of subunits forming the channel complex may simply reflect clustering of several ENaC channels at the cell surface.

Structure and Function Relationships

The Channel Pore

The recognition of amino acid residues on α -, β -, and γ -ENaC subunits that when mutated drastically change the channel affinity for the pore blocker amiloride (up to a 1000-fold increase in amiloride K_i) (36), identified a sequence (α S583,

β G525, and γ G537 in the rat ENaC sequences) that likely forms the outer vestibule of the channel where amiloride binds (see Figure 2). These amino acids are located at the extracellular start of the second transmembrane α helix; mutations within this region affect the amiloride-blocking kinetics, indicating that the mutated amino acids participate in the close interaction between amiloride and the channel pore, *i.e.*, are part of the amiloride binding site (37).

It is likely that other regions in the ENaC subunit sequence also influence channel affinity for amiloride, but these regions remain to be identified. For instance, mutations in the WYR-FHY sequence lower the affinity for amiloride of a 21-pS channel made of α -ENaC subunits exclusively (38,39). There is no evidence that these mutations affect the block by amiloride of the native 5-pS channel made of $\alpha\beta\gamma$ subunits and thus participate in the amiloride binding site in the native channel.

According to an early model of the ENaC pore based on the interaction between pore blockers, large blocking cations and the permeating Na^+ and Li^+ ions, it was proposed that the external vestibule of ENaC resembles a funnel-like structure that narrows from the amiloride binding site down to the selectivity filter, allowing only small permeant ions to pass through the channel pore (40). Consistent with this model, mutations of conserved Gly and Ser residues downstream of the amiloride binding site in the rat α -ENaC sequence (GSS sequence) and located within the first five amino acid residues of the second transmembrane α helix change the permeability properties of the channel (41–43). First, these mutations allow larger cations to pass through the channel but also reduce the unitary conductance of ENaC, *i.e.*, the flux of Na^+ or Li^+ ions through the pore. These experiments indicate that these mutations disrupt the channel structure at a site where the permeating ions are in close contact with the channel pore, likely the selectivity filter.

Thus, the stretch of seven amino acid residues at the external start of the transmembrane α helix that compose the amiloride binding site and the selectivity filter line the external entry of the channel pore. The structures that line the internal pore and its opening into the cytosol remain to be identified.

Channel Gates

Secondary structure predictions reveal a short external α helix (pore helix) upstream of the second transmembrane segment (Figure 2). Within this external vestibule helix, a mutation of a conserved Ala residue in degenerin channels causes degeneration of touch receptor cells in *Caenorhabditis elegans* with morphologic features that are consistent with an abnormal cation leak into the cell (44). At the corresponding position in α -ENaC, introduction of a cysteine for a conserved Ser (α ENaC S576) and its covalent modification with sulfhydryl reagent induces changes in the ENaC gating with extremely long channel opening dwell times ($P_o > 0.9$) (45,46). It is interesting that the Cys at this position 576 is accessible by the sulfhydryl reagent only when the channel is open. These observations are consistent with the presence of an extracellular gate controlling the channel openings. The putative external vestibule helix may function as an external gate and the sulf-




Figure 2. Schematic representation of the ENaC channel in the membrane. (Left) Secondary structure predictions and membrane topology of one ENaC subunit. Each subunit is made up of two transmembrane segments, TM1 and TM2. Cylinders represent putative α helices. P, extracellular pore helix; HG, conserved His and Gly residues among all of the members of the ENaC/degenerins family members; P.Y, PY motif common to the $\alpha\beta\gamma$ -ENaC subunits; DEG, degenerin site in the P-helix. (Right) ENaC channel model. The external pore vestibule is made up of the P- α helix, the linker between P and TM2 helix that composes the amiloride binding site. The first residues of TM2 participate in the ion selectivity filter that presents the narrowest part of the ion conduction pathway, where the permeant ion is in its dehydrated form. Structures that line the internal water-filled pore remain to be identified. Illustration by Josh Gramling—Gramling Medical Illustration.

hydriyl reagent as a "foot in the door" that locks the channel in the open conformation.

Evidence for an intracellular gating domain first came from genetic analysis of ENaC loss-of-function mutations associated with pseudohypoaldosteronism type 1. It was found that in one family, the disease results from a mutation of a highly conserved His-Gly motif in the N-terminus of the β -ENaC subunit (47). Corresponding mutations in the α -ENaC subunit modifies channel gating by shortening the channel mean open time and therefore decreasing the Po (48). This His-Gly motif that is conserved among all of the members of the ENaC/degenerin channel family is located in an intracellular linker between a putative hydrophobic α helix and the first transmembrane segment. Furthermore, this region of the N-terminus in the $\alpha\beta\gamma$ -ENaC subunits is rich in cysteine residues. These residues are responsible for the high ENaC sensitivity to inhibition by a variety of intracellular sulfhydryl reagents, including methanethiosulfonates, metal divalent cations, and oxidizing agents (49). These reagents inhibit ENaC activity from the cytosolic side by inducing long and slowly reversible channel closures. The contribution of the cytosolic C-terminus of $\alpha\beta\gamma$ -ENaC subunits in channel gating still needs to be elucidated.

Modulatory Sites of ENaC Activity

Aldosterone and many other factors, including hormones (e.g., vasopressin, angiotensin II, insulin), extracellular proteases (e.g., kallikrein, channel-activating protease-1), intra- and extracellular ion concentrations (e.g., for Na^+ , Ca^{2+}), osmolarity, and the tubular flow rate are known to regulate ENaC activity (1). The regulation of renal sodium transport by ENaC may occur directly either at the level of the single channel by changing the Po present in the apical membrane or at the level of the number (N) of active channels expressed at the cell surface by changing the rate of insertion or retrieval from the cell surface. These two ways of regulation of ENaC activity at the cell surface are not mutually exclusive. In the case of aldosterone-dependent stimulation of Na^+ transport, the insertion of additional ENaC subunits into the apical membrane is supported by immunohistochemical studies (11,12) and single-channel recordings show that ENaC are more active at the cell surface (20,24). Aldosterone regulation of ENaC likely involves both effects on N and Po.

The SGK1 kinase, a member of the PKB/Akt family of serine/threonine kinases, is rapidly induced by aldosterone in ASDN model epithelia *in vitro* (50,51) as well as in the kidney *in vivo* (52,53). Co-expression of ENaC with SGK1 in heterologous expression systems profoundly increases ENaC-mediated Na^+ currents (50,51,54) and ENaC cell surface expression. The regulation of ENaC by SGK can be mediated either directly by phosphorylation of targeted amino acid residues in the ENaC sequence (55) or indirectly by phosphorylating ENaC regulatory proteins. In heterologous expression systems, the effect of SGK1 on ENaC can be mediated by phosphorylation of the ubiquitin-ligase *Nedd4-2* that regulates ENaC activity at the cell surface (56,57).

Nedd4-2 is a ubiquitin ligase that contains in its sequence four WW domains (W for tryptophane) that bind proline-rich

ligands. Proline-rich motifs are found in the C-terminus of the $\alpha\beta\gamma$ -ENaC subunits, in particular a PP.Y sequence (PY motif) that represents a characteristic pattern for WW peptide ligands. These PPP.Y sequences in the β - and γ -ENaC subunits are the target sequence for mutations that cause Liddle's syndrome (pseudohypoaldosteronism) (58–60). These mutations increase the number of channel molecules at the cell surface as well as the channel activity (Po effect) when expressed in *Xenopus* oocytes (61). Mutations of the PY motif clearly prolong the half-life of the channel at the cell surface as a result of an impaired internalization of ENaC. Two mechanisms have been proposed to explain the cell surface retention of ENaC: First, mutations in the PY motif impair interactions with the binding partner *Nedd4-2*; because *Nedd4-2* catalyzes the attachment of ubiquitin moieties on the channel for endocytosis and degradation, mutations in the PY motif alter this process (62). Second is a defect in the clathrin-mediated endocytosis of ENaC, because the PY motif sequence also contains two adjacent endocytic signals (63). Although these proposed mechanisms are not mutually exclusive, it remains that in heterologous expression systems, ENaC is ubiquitinated, and multiple lysine residues in the N-termini of α (K47 and K50) and γ subunits (K6 to K13) are substrates for attachment of ubiquitin moieties (64). How mutations in the PY motif increase the Po remains to be established. There is presently no good evidence that the C-terminus of ENaC subunits directly participates in channel gating, but it remains possible that the protein interaction with the PY motif of ENaC downregulates the channel (Po effect) before internalization.

Because the PY motif in the C-terminus of ENaC subunits regulates ENaC stability and activity at the cell surface in heterologous expression systems, studies have addressed the question of whether this PY motif represents a final target sequence on ENaC for the aldosterone signaling pathway. In their early report on patients with pseudohypoaldosteronism, Liddle *et al.* (65) clearly showed that these patients retain their ability to respond to aldosterone by reducing their urinary fractional excretion of sodium to almost zero. More recently, single-channel studies in mouse models of Liddle syndrome (pseudohypoaldosteronism) that lack the PY motif in the β -ENaC subunit show that the ENaC response to elevation in plasma aldosterone level is drastically increased in Liddle mice compared with wild-type mice (66). In a cortical CD cell line that expressed ENaC wild-type or mutants with mutations in the PY motif, the rate of increase in Na^+ transport induced by aldosterone during the early response was similar (67). These observations suggest that the PY motif regulates ENaC activity at the cell surface independent of a stimulation of the aldosterone signaling pathway.

Regarding the possibility that SGK1 directly phosphorylates ENaC, the cytosolic C-terminus of the mammalian α -ENaC subunit contains consensus sites for phosphorylation by SGK. The mutation of the serine 621 (rENaC sequence) prevents the effect of SGK1 that recruits more active channel at the cell surface, suggesting that S621 of α -ENaC also participates in the control of ENaC expression at the cell surface (55).

Extracellular trypsin and proteases CAP1 (prostasin), CAP2,

CAP3, and TMPRSS3 activate ENaC (68,69). In addition, the serine protease inhibitor aprotinin reduces the transepithelial sodium transport in an amphibian kidney cell line. In airway epithelial cell lines, elastase, another serine protease, activates ENaC, and a decrease in the expression of prostasin reduces the amiloride-sensitive Na^+ current (70,71). Clearly, the Po of ENaC is increased by extracellular proteases, an effect on the channel that is independent of changes in intracellular Ca^{2+} or activation of G protein-coupled receptors (72,73). In addition, maturation of ENaC involves proteolytic cleavage of the α - and γ -ENaC subunits by furin, a proprotein convertase that also cleaves prohormones, coagulation factors, or receptors at the cell surface (74). The furin-dependent cleavage of ENaC seems to correlate with a higher channel activity at the cell surface, and the relation between ENaC cleavage and the activation of the channel was confirmed in furin-deficient cells (74).

ENaC contains several consensus motifs for furin or trypsin cleavage in the extracellular loop. Furin cleavage of α - and γ -ENaC involves multiple cleavage motifs in the extracellular loop after the first transmembrane segment, suggesting the presence in this region of important functional domains that control ENaC activity (74). These cleavage sites do not overlap with sites for trypsin or serine protease action, which remain to be precisely identified.

Other modulatory sites are likely located in the extracellular loop. For instance, mutations of histidine residues at corresponding positions in the extracellular loop of α - and γ -ENaC differentially modulate ENaC self-inhibition by extracellular Na^+ ions (75). Thus, the extracellular loop may function as an extracellular sensor for Na^+ ions to adapt channel activity with the extracellular concentration of Na^+ ions; in addition, the extracellular loop of ENaC may function as a target for an extracellular signaling cascade involving proteases, necessary for the activation of the channel.

Conclusion

The identification of structural domains that are important for basic channel function (e.g., translocation of Na^+ ions across the membrane) or for the modulation of ENaC activity at the cell surface (N or Po effects) is essential for our understanding of the cellular and molecular mechanisms involved in ENaC regulation by hormones and other intracellular/extracellular factors. In addition, this knowledge about the structure and the function of ENaC will certainly be helpful in interpreting physiologic and pathophysiologic consequences of ENaC genetic variants on Na^+ handling by the ASDN.

Acknowledgments

The authors' work is supported by the Swiss National Science Foundation (to J.L., grant no. 3200B0-10579/1; to L.S., grant no. 3100A0-108069/1). We wish to apologize to those authors whose work could not be cited due to space restrictions.

References

1. Kellenberger S, Schild L: Epithelial sodium channel/de-

- generin family of ion channels: A variety of functions for a shared structure. *Physiol Rev* 82: 735–767, 2002
2. Rossier BC, Pradervand S, Schild L, Hummler E: Epithelial sodium channel and the control of sodium balance: Interaction between genetic and environmental factors. *Annu Rev Physiol* 64: 877–897, 2002
3. Oh YS, Warnock DG: Disorders of the epithelial Na^+ channel in Liddle's syndrome and autosomal recessive pseudohypoaldosteronism type 1. *Exp Nephrol* 8: 320–325, 2000
4. Reilly RF, Ellison DH: Mammalian distal tubule: Physiology, pathophysiology, and molecular anatomy. *Physiol Rev* 80: 277–313, 2000
5. Bachmann S, Bostanjoglo M, Schmitt R, Ellison DH: Sodium transport-related proteins in the mammalian distal nephron—Distribution, ontogeny and functional aspects. *Anat Embryol (Berl)* 200: 447–468, 1999
6. Loffing J, Kaissling B: Sodium and calcium transport pathways along the mammalian distal nephron: From rabbit to human. *Am J Physiol Renal Physiol* 284: F628–F643, 2003
7. Bostanjoglo M, Reeves WB, Reilly RF, Velazquez H, Robertson N, Litwack G, Morsing P, Dorup J, Bachmann S, Ellison DH: 11 β -hydroxysteroid dehydrogenase, mineralocorticoid receptor, and thiazide-sensitive Na-Cl cotransporter expression by distal tubules. *J Am Soc Nephrol* 9: 1347–1358, 1998
8. Masilamani S, Wang XY, Kim GH, Brooks H, Nielsen J, Nielsen S, Nakamura K, Stokes JB, Knepper MA: Time course of renal Na-K-ATPase, NHE3, NKCC2, NCC, and ENaC abundance changes with dietary NaCl restriction. *Am J Physiol Renal Physiol* 283: F648–F657, 2002
9. Kim GH, Masilamani S, Turner R, Mitchell C, Wade JB, Knepper MA: The thiazide-sensitive Na-Cl cotransporter is an aldosterone-induced protein. *Proc Natl Acad Sci U S A* 95: 14552–14557, 1998
10. Velazquez H, Bartiss A, Bernstein P, Ellison DH: Adrenal steroids stimulate thiazide-sensitive NaCl transport by rat renal distal tubules. *Am J Physiol* 270: F211–F219, 1996
11. Masilamani S, Kim GH, Mitchell C, Wade JB, Knepper MA: Aldosterone-mediated regulation of ENaC α , β , and γ subunit proteins in rat kidney. *J Clin Invest* 104: R19–R23, 1999
12. Loffing J, Pietri L, Aregger F, Bloch-Faure M, Ziegler U, Meneton P, Rossier BC, Kaissling B: Differential subcellular localization of ENaC subunits in mouse kidney in response to high- and low-Na diets. *Am J Physiol Renal Physiol* 279: F252–F258, 2000
13. Frindt G, Palmer LG: Na channels in the rat connecting tubule. *Am J Physiol Renal Physiol* 286: F669–F674, 2004
14. Costanzo LA: Comparison of calcium and sodium transport in early and late rat distal tubules: Effect of amiloride. *Am J Physiol* 246: F937–F945, 1984
15. Tomita K, Pisano JJ, Knepper MA: Control of sodium and potassium transport in the cortical collecting duct of the rat. Effects of bradykinin, vasopressin, and deoxycorticosterone. *J Clin Invest* 76: 132–136, 1985
16. Almeida AJ, Burg MB: Sodium transport in the rabbit connecting tubule. *Am J Physiol* 243: F330–F334, 1982
17. Katz AI, Doucet A, Morel F: Na-K-ATPase activity along the rabbit, rat, and mouse nephron. *Am J Physiol* 237: F114–F120, 1979
18. Rubera I, Loffing J, Palmer LG, Frindt G, Fowler-Jaeger N,

- Sauter D, Carroll T, McMahon A, Hummler E, Rossier BC: Collecting duct-specific gene inactivation of alphaENaC in the mouse kidney does not impair sodium and potassium balance. *J Clin Invest* 112: 554–565, 2003
19. Verrey F, Hummler E, Schild L, Rossier BC: Control of Na⁺ transport by aldosterone. In: *The Kidney*, 3rd Ed., edited by Seldin DW, Giebisch G, Philadelphia, Lippincott Williams & Wilkins, 2001, pp 1441–1472
 20. Hamilton KL, Eaton DC: Single-channel recordings from amiloride-sensitive epithelial sodium channel. *Am J Physiol* 249: C200–C207, 1985
 21. Palmer LG, Frindt G: Amiloride-sensitive Na channels from the apical membrane of the rat cortical collecting tubule. *Proc Natl Acad Sci U S A* 83: 2727–2770, 1986
 22. Palmer LG, Frindt G: Gating of Na channels in the rat cortical collecting tubule: Effects of voltage and membrane stretch. *J Gen Physiol* 107: 35–45, 1996
 23. Palmer LG, Frindt G: Effects of cell Ca and pH on Na channels from rat cortical collecting tubule. *Am J Physiol* 253: F333–F339, 1987
 24. Frindt G, Masilamani S, Knepper MA, Palmer LG: Activation of epithelial Na channels during short-term Na deprivation. *Am J Physiol Renal Physiol* 280: F112–F118, 2001
 25. Chraïbi A, Horisberger AD: Na self inhibition of human epithelial Na channel: Temperature dependence and effect of extracellular proteases. *J Gen Physiol* 120: 133–145, 2002
 26. Palmer LG, Andersen OS: Interactions of amiloride and small monovalent cations with the epithelial sodium channel. Inferences about the nature of the channel pore. *Biophys J* 55: 779–787, 1989
 27. Palmer LG: Interactions of amiloride and other blocking cations with the apical Na channel in the toad urinary bladder. *J Membr Biol* 87: 191–199, 1985
 28. Burch LH, Talbot CR, Knowles MR, Canessa CM, Rossier BC, Boucher RC: Relative expression of the human epithelial Na⁺ channel subunits in normal and cystic fibrosis airways. *Am J Physiol* 269: C511–C518, 1995
 29. Canessa CM, Schild L, Buell G, Thorens B, Gautschi I, Horisberger J-D, Rossier BC: Amiloride-sensitive epithelial Na⁺ channel is made of three homologous subunits. *Nature* 367: 463–467, 1994
 30. Canessa CM, Merillat A-M, Rossier BC: Membrane topology of the epithelial sodium channel in intact cells. *Am J Physiol Cell Physiol* 267: C1682–C1690, 1994
 31. Firsov D, Gautschi I, Merillat AM, Rossier BC, Schild L: The heterotetrameric architecture of the epithelial sodium channel (ENaC). *EMBO J* 17: 344–352, 1998
 32. Coscoy S, Lingueglia E, Lazdunski M, Barbry P: The pHEMT-arg-pHE-amide-activated sodium channel is a tetramer. *J Biol Chem* 273: 8317–8322, 1998
 33. Kosari F, Sheng SH, Li JQ, Mak DD, Foscett JK, Kleyman TR: Subunit stoichiometry of the epithelial sodium channel. *J Biol Chem* 273: 13469–13474, 1998
 34. Dijkink L, Hartog A, Van Os CH, Bindels RJM: The epithelial sodium channel (ENaC) is intracellularly located as a tetramer. *Pflugers Arch* 444: 549–555, 2002
 35. Snyder PM, Cheng C, Prince LS, Rogers JC, Welsh MJ: Electrophysiological and biochemical evidence that deg/enac cation channels are composed of nine subunits. *J Biol Chem* 273: 681–684, 1998
 36. Schild L, Schneeberger E, Gautschi I, Firsov D: Identification of amino acid residues in the alpha, beta, gamma subunits of the epithelial sodium channel (ENaC) involved in amiloride block and ion permeation. *J Gen Physiol* 109: 15–26, 1997
 37. Kellenberger S, Gautschi I, Schild L: Mutations in the epithelial Na⁺ channel ENaC outer pore disrupt amiloride block by increasing its dissociation rate. *Mol Pharmacol* 64: 848–856, 2003
 38. Ismailov II, Kieberemmons T, Lin CM, Berdiev BK, Shlyonsky VG, Patton HK, Fuller CM, Worrell R, Zuckerman JB, Sun W, Eaton DC, Benos DJ, Kleyman TR: Identification of an amiloride binding domain within the alpha-subunit of the epithelial Na⁺ channel. *J Biol Chem* 272: 21075–21083, 1997
 39. Kelly O, Lin CM, Ramkumar M, Saxena NC, Kleyman TR, Eaton DC: Characterization of an amiloride binding region in the alpha-subunit of ENaC. *Am J Physiol Renal Physiol* 285: F1279–F1290, 2003
 40. Palmer LG: Epithelial Na channels: The nature of the conducting pore. *Renal Physiol Biochem* 13: 51–58, 1990
 41. Kellenberger S, Hoffmann-Pochon N, Gautschi I, Schneeberger E, Schild L: On the molecular basis of ion permeation in the epithelial Na⁺ channel. *J Gen Physiol* 114: 13–30, 1999
 42. Kellenberger S, Auberson M, Gautschi I, Schneeberger E, Schild L: Permeability properties of ENaC selectivity filter mutants. *J Gen Physiol* 118: 679–692, 2001
 43. Kellenberger S, Gautschi I, Schild L: A single point mutation in the pore region of the epithelial Na⁺ channel changes ion selectivity by modifying molecular sieving. *Proc Natl Acad Sci U S A* 96: 4170–4175, 1999
 44. Driscoll M, Chalfie M: The mec-4 gene is a member of a family of *Caenorhabditis elegans* genes that can mutate to induce neuronal degeneration. *Nature* 349: 588–593, 1991
 45. Snyder PM, Bucher DB, Olson DR: Gating induces a conformational change in the outer vestibule of ENaC. *J Gen Physiol* 116: 781–790, 2000
 46. Kellenberger S, Gautschi I, Schild L: An external site controls closing of the epithelial Na⁺ channel ENaC. *J Physiol (Lond)* 543: 413–424, 2002
 47. Grunder S, Firsov D, Chang SS, Jaeger NF, Gautschi I, Schild L, Lifton RP, Rossier BC: A mutation causing pseudohypoaldosteronism type 1 identifies a conserved glycine that is involved in the gating of the epithelial sodium channel. *EMBO J* 16: 899–907, 1997
 48. Grunder S, Jaeger NF, Gautschi I, Schild L, Rossier BC: Identification of a highly conserved sequence at the N-terminus of the epithelial Na⁺ channel alpha subunit involved in gating. *Pflugers Arch* 438: 709–715, 1999
 49. Kellenberger S, Gautschi I, Pfister Y, Schild L: Intracellular thiol-mediated modulation of ENaC activity. *J Biol Chem* 280: 7739–7747, 2005
 50. Naray FT, Canessa C, Cleaveland ES, Aldrich G, Fejes-Toth G: SgK is an aldosterone-induced kinase in the renal collecting duct—Effects on epithelial Na⁺ channels. *J Biol Chem* 274: 16973–16978, 1999
 51. Chen SY, Bhargava A, Mastroberardino L, Meijer OC, Wang J, Buse P, Firestone GL, Verrey F, Pearce D: Epithelial sodium channel regulated by aldosterone-induced protein sgk. *Proc Natl Acad Sci U S A* 96: 2514–2519, 1999
 52. Loffing J, Zecevic M, Feraille E, Kaissling B, Asher C, Rossier BC, Firestone GL, Pearce D, Verrey F: Aldosterone induces rapid apical translocation of ENaC in early portion

- of renal collecting system: Possible role of SGK. *Am J Physiol Renal Physiol* 280: F675–F682, 2001
53. Muller OG, Parnova RG, Centeno G, Rossier BC, Firsov D, Horisberger JD: Mineralocorticoid effects in the kidney: Correlation between α ENaC, GILZ, and Sgk-1 mRNA expression and urinary excretion of Na^+ and K^+ . *J Am Soc Nephrol* 14: 1107–1115, 2003
 54. Alvarez de la Rosa D, Zhang P, Naray-Fejes-Toth A, Fejes-Toth G, Canessa CM: The serum and glucocorticoid kinase sgk increases the abundance of epithelial sodium channels in the plasma membrane of *Xenopus* oocytes. *J Biol Chem* 274: 37834–37839, 1999
 55. Diakov A, Korbmacher C: A novel pathway of epithelial sodium channel activation involves a serum- and glucocorticoid-inducible kinase consensus motif in the C terminus of the channel's α -subunit. *J Biol Chem* 279: 38134–38142, 2004
 56. Debonneville C, Flores SY, Kamynina E, Plant PJ, Tauxe C, Thomas MA, Munster C, Chraïbi A, Pratt JH, Horisberger JD, Pearce D, Loffing J, Staub O: Phosphorylation of Nedd4-2 by Sgk1 regulates epithelial Na^+ channel cell surface expression. *EMBO J* 20: 7052–7059, 2001
 57. Snyder PM, Olson DR, Thomas BC: Serum and glucocorticoid-regulated kinase modulates Nedd4-2-mediated inhibition of the epithelial Na^+ channel. *J Biol Chem* 277: 5–8, 2002
 58. Shimkets RA, Warnock DG, Bositis CM, Nelson-Williams C, Hansson JH, Schambelan M, Gill JR Jr, Ulick S, Milora RV, Findling JW, Canessa CM, Rossier BC, Lifton RP: Liddle's syndrome: Heritable human hypertension caused by mutations in the β subunit of the epithelial sodium channel. *Cell* 79: 407–414, 1994
 59. Hansson JH, Nelson-Williams C, Suzuki H, Schild L, Shimkets R, Lu Y, Canessa C, Iwasaki T, Rossier B, Lifton RP: Hypertension caused by a truncated epithelial sodium channel γ subunit: Genetic heterogeneity of Liddle syndrome. *Nat Genet* 11: 76–82, 1995
 60. Hansson JH, Schild L, Lu Y, Wilson TA, Gautschi I, Shimkets R, Nelson-Williams C, Rossier BC, Lifton RP: A *de novo* missense mutation of the β subunit of the epithelial sodium channel causes hypertension and Liddle syndrome, identifying a proline-rich segment critical for regulation of channel activity. *Proc Natl Acad Sci U S A* 92: 11495–11499, 1995
 61. Firsov D, Schild L, Gautschi I, M  rillat A-M, Schneeberger E, Rossier BC: Cell surface expression of the epithelial Na^+ channel and a mutant causing Liddle syndrome: A quantitative approach. *Proc Natl Acad Sci U S A* 93: 15370–15375, 1996
 62. Staub O, Dho S, Henry PC, Correa J, Ishikawa T, McGlade J, Rotin D: WW domains of Nedd4 bind to the proline-rich PY motifs in the epithelial Na^+ channel deleted in Liddle's syndrome. *EMBO J* 15: 2371–2380, 1996
 63. Shimkets RA, Lifton RP, Canessa CM: The activity of the epithelial sodium channel is regulated by clathrin-mediated endocytosis. *J Biol Chem* 272: 25537–25541, 1997
 64. Staub O, Abriel H, Plant P, Ishikawa T, Kanelis V, Saleki R, Horisberger JD, Schild L, Rotin D: Regulation of the epithelial Na^+ channel by Nedd4 and ubiquitination. *Kidney Int* 57: 809–815, 2000
 65. Liddle GW, Bledsoe T, Coppage WS: A familial renal disorder simulating primary aldosteronism but with negligible aldosterone secretion. *Trans Assoc Am Physicians* 76: 199–213, 1963
 66. Dahlmann A, Pradervand S, Hummler E, Rossier BC, Frindt G, Palmer LG: Mineralocorticoid regulation of epithelial Na^+ channels is maintained in a mouse model of Liddle's syndrome. *Am J Physiol Renal Physiol* 285: F310–F318, 2003
 67. Auberson M, Hoffmann-Pochon N, Vandewalle A, Kellenberger S, Schild L: Epithelial Na^+ channel mutants causing Liddle's syndrome retain ability to respond to aldosterone and vasopressin. *Am J Physiol Renal Physiol* 285: F459–F471, 2003
 68. Vuagniaux G, Vallet V, Jaeger NF, Pfister C, Bens M, Farman N, Courtois-Couty N, Vandewalle A, Rossier BC, Hummler E: Activation of the amiloride-sensitive epithelial sodium channel by the serine protease mCAP1 expressed in a mouse cortical collecting duct cell line. *J Am Soc Nephrol* 11: 828–834, 2000
 69. Vallet V, Chraïbi A, Gaeggeler HP, Horisberger JD, Rossier BC: An epithelial serine protease activates the amiloride-sensitive sodium channel. *Nature* 389: 607–610, 1997
 70. Caldwell RA, Boucher RC, Stutts MJ: Neutrophil elastase activates near-silent epithelial Na^+ channels and increases airway epithelial Na^+ transport. *Am J Physiol Lung Cell Mol Physiol* 288: L813–L819, 2005
 71. Tong Z, Illek B, Bhagwandin VJ, Verghese GM, Caughey GH: Prostaticin, a membrane-anchored serine peptidase, regulates sodium currents in JME/CF15 cells, a cystic fibrosis airway epithelial cell line. *Am J Physiol Lung Cell Mol Physiol* 287: L928–L935, 2004
 72. Chraïbi A, Vallet V, Firsov D, Hess SK, Horisberger JD: Protease modulation of the activity of the epithelial sodium channel expressed in *Xenopus* oocytes. *J Gen Physiol* 111: 127–138, 1998
 73. Caldwell RA, Boucher RC, Stutts MJ: Serine protease activation of near-silent epithelial Na^+ channels. *Am J Physiol Cell Physiol* 286: C190–C194, 2004
 74. Hughey RP, Bruns JB, Kinlough CL, Harkleroad KL, Tong QS, Carattino MD, Johnson JP, Stockand JC, Kleyman TR: Epithelial sodium channels are activated by furin-dependent proteolysis. *J Biol Chem* 279: 18111–18114, 2004
 75. Sheng S, Bruns JB, Kleyman TR: Extracellular histidine residues crucial for Na^+ self-inhibition of epithelial Na^+ channels. *J Biol Chem* 279: 9743–9749, 2004